

PRODUCT SPECIFICATIONS

Date: 2015-09-28

PRODUCT NAME	:	Anti-h cTnI 9708 SPTN-5
PRODUCT SPECIFICITY	:	Antibody recognizes human Cardiac Troponin I
PRODUCT CODE	:	100523
PRODUCT BUFFER	:	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
SHELF LIFE AND STORAGE	:	Shelf life unspecified, storage at 2-8 °C
ANALYTE DESCRIPTION	:	Troponins form a complex (Troponin C, I and T) of three regulatory proteins that are integral to muscle contraction in skeletal and cardiac muscle, but not smooth muscle. Troponin levels can be used as a test of several different heart disorders, including myocardial infarction. The measurement of serum cTnI and cTnT is superior in terms of sensitivity and specificity to cardiac muscle enzyme measurements in the identification of cardiac muscle damage. Raised cardiac troponin concentrations are now accepted as the standard biochemical marker for the diagnosis of myocardial infarction.

PARAMETERS TESTED FROM EACH LOT

PRODUCT APPEARANCE	:	Clear liquid, may turn opaque during storage.
PRODUCT CONCENTRATION	:	5.0 mg/ml (+/- 10 %)
PRODUCT ACTIVITY	:	80-120 % compared to reference in an IFMA-test
IEF-RANGE	:	5.9 – 6.5
PURITY	:	≥ 95 %

PARAMETERS DETERMINED ONLY DURING PRODUCT R&D PHASE

CLASS AND SUBCLASS	:	IgG ₁
ASSOCIATION CONSTANT	:	N/D
DISSOCIATION CONSTANT	:	N/D
AFFINITY CONSTANT	:	N/D
DETERMINATION METHOD	:	-
ANTIGEN	:	-
CROSS-REACTIVITIES	:	Does not react with skeletal troponin I. Others not tested.

EPITOPE : N-terminal end of the molecule (residues 22-43).

EPITOPE GROUP :

-
Two antibodies binding to the same, or closely located epitopes, belong to the same group and hence cannot be used as a pair in a sandwich assay. Epitope group numbering does not give any detailed information where the epitope is located.

PAIR RECOMMENDATIONS :

SOLID	LABEL
-	-

Please note that pair recommendations are based on results obtained in our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations should be taken only as a directive.

PRODUCT STABILITY :

TEMPERATURE, DAYS	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 7 days	OK
+35 °C, 21 days	OK, but charge changes
+45 °C, 3 days	OK
+45 °C, 7 days	OK, but charge changes

Please note that the shelf life given on page one is based on real time stability testing at +2-8 °C in the product buffer.

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. The maximum duration of the test is 21 days, except for the +45 °C only 7 days.

MISCELLANEOUS :

-

REFERENCES :

-

Legal disclaimer

MedixMAB monoclonal antibodies meet their specifications if transported, stored and used according to the instructions for use. See also separate *Notes on use and storage of monoclonal antibodies*. If not otherwise agreed in writing all products are sold under *General Delivery Terms*. Other terms may apply if MedixMAB monoclonal antibodies are purchased from an official distributor of Medix Biochemica products. MedixMAB is a registered trademark of Medix Biochemica and may not be used or reproduced without Medix Biochemica's written permission. Further information on products and methods is available from medixMAB website.